

BUSTER

Contain:Bacillus thuringiensis 1.5x10⁸cfu/ml, vegetable oil 2% (glycerol-inert material)

Fertilizer

Formulation:soluble concentrate

Hazard phrases

Not applicable

Precautionary phrases

P501 - Dispose of contents/container to an authorized waste disposal station.

it is IT

Lot no.:see above

Date of manufacture:see above

Expiration date:24 months

Storage:Store in a dry, cool and well-ventilated place..Storage temperature 5°C-25°C

Careful:Read the attached instructions before use.

RO-FERTILIZER Authorization No.

Producer:Norofert SA

LT. Av. Serban Petrescu Street, No. 20, Sector 1,

Bucharest, Romania

Phone: 0766 080 767

For your attention:Shake the container vigorously for 15-20 seconds for good homogenization. By following the instructions presented, the product can be used for its intended purpose. The manufacturer declines any responsibility for damage due to storage, handling or use not in accordance with the recommendations on this label.

BUSTER

Mode of action:

Beneficial effect of *B. thuringiensis* as a biofertilizer and plant growth promoter. Promotes growth, tolerance to drought stress and increases nutrient uptake and utilization. Can lead to increased stem height, root length, fruit weight and total dry weight in plants. Plants increase nutrient uptake in drought environments and increase macro and micronutrient intake. and helps plants survive environmental stresses such as heavy metals. Once in the soil it has the ability to convert insoluble P into soluble form.

How to use:

Culture	Dose
Sunflower	1 l/ha
Winter wheat	1 l/ha
Autumn rapeseed	1 l/ha
Maize	1 l/ha
SOYBEAN	1 l/ha

Application time:

It can be applied throughout the growing season, in multiple applications at 14-21 day intervals.

Preparation of the solution:

Fill the sprayer tank halfway, add the appropriate amount of product, and then top up with water. Stir the spray solution well during filling and application. Use clean water from a known source.

Application technique:

Ground equipment can be used for field application of the products, commonly used on the above-mentioned crops. The volume of solution used depends on the equipment used. It is recommended to use a solution volume of 200-400 l/ha, so that there is good penetration and coverage of the crop. The solution will be applied immediately after mixing.

Application recommendations:

Do not store the prepared solution overnight. Avoid applying in the middle of the day when temperatures exceed 25°C. If rain occurs shortly after application, it will be necessary to repeat the treatment. Do not apply when wind speed favors drift beyond the area intended for treatment.

Compatibility: Mixing with other products should be avoided to prevent inactivation of microorganisms.

Break time: 0 days

Cleaning the equipment: Equipment will be pressure washed before and after application, and particularly if a product that could damage the crop was used before Buster.